

***Peltigera neodegenii* sp. nov. from Central China**

LIU-FU HAN^{1#}, XIAO-MIN XU^{1#},
JING-YUAN YANG^{2#}, SHOU-YU GUO^{3*}

¹ College of Life Science, Hebei Normal University,
Shijiazhuang 050024, P. R. China

² Shennongjia National Nature Reserve Administration,
Shennongjia, Hubei 442400, P. R. China

³ State Key Laboratory of Mycology, Institute of Microbiology,
Chinese Academy of Sciences, Beijing 100101, P. R. China

* CORRESPONDENCE TO: guosy@im.ac.cn

ABSTRACT—*Peltigera neodegenii* from Central China is described and illustrated as a new species. It is similar in general appearance to *P. degenii*, *P. membranacea*, and *P. canina*, but is distinguished by its shiny upper surface in the central part of lobes, tomenta around the margins of upper surface of lobes, distinct and raised veins on the lower surface of lobes, and simple rhizines. Comparisons of the ITS (ITS1, 5.8S, ITS2) sequences of the nuclear ribosomal DNA repeat tandem, both in phylogenetic analysis and secondary structure models of ITS2, support the taxonomic distinctness of this species.

KEY WORDS—biodiversity, cyanolichen, *Peltigeraceae*, *Peltigerales*, Shennongjia Mountain

Introduction

Peltigera Willd. (*Peltigeraceae*), characterized by large foliose thalli and a worldwide distribution in moist habitats, currently includes more than 90 species (Kirk & al. 2008; Han & al. 2013, 2015; Manoharan-Basil & al. 2016). In modern lichen taxonomy, molecular techniques have been used widely to delimit species within the genus (Goffinet & Miądlikowska 1999, Sérusiaux & al. 2009). The nrDNA ITS region, the main molecular marker or barcode used to characterize fungal and lichen species (Schoch & al. 2012, Guo 2013),

LIU-FU HAN, XIAO-MIN XU & JING-YUAN YANG contributed equally to this work.

has been helpful in revealing phylogenetic relationships in *Peltigera* (Goffinet & al. 2003; O'Brien & al. 2009; Sérusiaux & al. 2009; Han & al. 2013, 2015; Manoharan-Basil & al. 2016).

Peltigera has been relatively well studied in the northern hemisphere and a phylogeny combining morphological and chemical characters with LSU nrDNA sequence analysis has been carried out (Miądlikowska & Lutzoni 2000). In Asia, about 40 species have been reported, and the nrDNA ITS sequence data of most species are available from GenBank (Martínez & al. 2003; Han & al. 2013, 2015). Of the 30 species recorded in China (Chen 1986; Vitikainen 1986; Wei 1991; Stenroos & al. 1994; Wu & Liu 2012; Han & al. 2013, 2015; Niu & al. 2016), 14 belong to the *P. canina* group, in *P.* sect. *Peltigera*. During our ongoing study of *Peltigera* species in China, we identified a new species based on morphological characteristics and nrDNA ITS sequence data, which we describe here as *Peltigera neodegenii*. We also compare the secondary structure of the *P. neodegenii* nrDNA ITS2 region with those of two closely related species, *P. degenii* Gyeln. and *P. membranacea* (Ach.) Nyl.

Materials & methods

Specimens & morphology

Specimens were collected from Shennongjia forest region in Central China and all examined morphologically using a Motic SMZ-140 or Zeiss-Stereo Discovery-V12 dissecting microscope and Leica DM500 compound microscope. Asci and ascospores were observed in sections of apothecia cut by hand with razor and mounted in water after staining in 0.2% toluidine blue for about 15 min. Descriptions and terminology follow Vitikainen (2004). Thin layer chromatography (TLC) was performed on all specimens using solvent systems C and G according to Orange & al. (2010). The specimens are deposited both in the Herbarium Mycologicum Academiae Sinicae-Lichenes, Beijing, China (HMAS-L) and the Lichen Section of Botanical Herbarium, Hebei Normal University, Shijiazhuang, China (HBNU).

DNA extraction, PCR amplification, sequencing

The samples of lobe tips were cut for DNA extraction from five specimens (including the type and three other *P. neodegenii* specimens and one *Peltigera praetextata* specimen). DNA was extracted using the DNAsecure Plant DNA Kit following the manufacturer's protocol. The ITS region was amplified according to Han & al. (2013, 2015). The entire nrITS region (ITS1+5.8S+ITS2) was targeted for Polymerase Chain Reaction (PCR) using primers ITS1F and ITS4 (White & al. 1990) in a 25 µL volume containing 0.75 units of TransStart Taq Polymerase, 2.5 µL of ITS buffer, 0.5 µL of a 5 µM solution of the primers, 2 µL of 2.5 mM for each dNTP solution, and 1 µL of genomic DNA. Thermal cycling conditions were 95°C for 3 min initiation, followed by 35 cycles at 94°C for 30 s, 54°C for 30 s, and 72°C for 1 min

with a final extension at 72°C for 10 min. PCR products were screened on 1% agarose gels stained with ethidium bromide and sequenced by the Genewiz Inc. (Suzhou, Jiangsu, China).

Phylogenetic analysis & sequence comparisons

The entire ITS sequences of five samples of our examined specimens and the 24 representatives selected from GenBank were aligned manually by both ClustalW and Muscle implemented in MEGA version6 (Tamura & al. 2013). We excluded the 3' end of the 18S gene and the 5' end of the 26S gene from the analyses. Sequences from the new species were aligned with GenBank ITS sequences from taxa sharing the most similarity with them as determined from morphological characters, Blast results of sequence data, and selected references (e.g., Miadlikowska & Lutzoni 2000, Goffinet & al. 2003, Miadlikowska & al. 2003, Sérusiaux & al. 2009).

Ambiguously aligned regions sensu Lutzoni & al. (2000) were delimited manually and excluded. The final matrix of the remaining 520 characters submitted to TreeBase with accession number TB2:S15264 may be obtained from the corresponding authors.

The evolutionary history was inferred from 29 nucleotide sequences by using both the Maximum Likelihood method based on the GTR+I model in MEGA6 and Bayesian inference (Huelsenbeck & Ronquist 2001) based on GTR model with rates = Invgamma. The secondary structures of ITS2 sequences for the new species and its most closely related species were predicted by The ITS2 Database III (Koetschan & al. 2010) and illustrated with PseudoViewer3 (Byun & Han 2009).

Results

Phylogenetic analysis and secondary structure of ITS2

The newly obtained sequences were submitted to GenBank. The entire ITS region, which was successfully sequenced for five samples from the collections from Mt. Shennongjia, Central China, comprises 573–574 bp (ITS1 = 216–217 bp; 5.8S = 157 bp; and ITS2 = 200 bp). We included ITS sequences from our new species and the GenBank reference sequences in the phylogenetic analyses. The data matrix of the aligned 29 sequences comprised 520 characters, of which 468 (90%) were constant and 52 (10%) were parsimony informative.

The hypervariable region in the internal transcribed spacer 1 (ITS1-HR; Miadlikowska & al. 2003) was calculated, with about 23% variable sites compared with *P. degenii*, for 86 positions in total (not shown).

The alignment dataset was analyzed using MrBayes for Bayesian inference and MEGA6 for Maximum Likelihood; similar topologies were obtained. A ML tree with bootstrap values (1000 replicats) and Bayesian posterior probabilities (BPP) at branches is shown in FIG. 1. In the phylogenetic tree, the new species formed a clade with *P. membranacea*. The different samples of the new species formed a clade with 70% ML support and 0.97 of BPP.




FIG. 1. Phylogenetic relationships among selected members (7 taxa, 29 sequences) closely related to *Peltigera neodegenii* based on ITS sequences. Values associated with internodes represent bootstrap support (ME-BP; before the slash) and posterior probability support (PP; after the slash). Only values $\geq 50\%$ for ML-BP and ≥ 0.90 for PP are shown. *Peltigera cinnamomea* Goward was included as outgroup. Fonts in bold indicate newly generated sequences from specimens collected in Central China; the ex-type sequence of *P. neodegenii* (MF085563) is annotated: T. There was a total of 520 positions in the final dataset. Evolutionary analyses were conducted in MEGA6 and MrBayes3.2.




FIG. 2. The secondary structure model for the ITS2 sequence of *Peltigera neodegenii* (right, GenBank MF085563 *P. neodegenii* ex-type) and its template G1255347364 (left, GenBank FJ709030 *P. degenii*). Red arrows indicate a hemi-compensatory base change (hCBC) and the green box indicates a compensatory base change (CBC).

A high-quality secondary structure model of the ITS2 sequence representing the new species (GenBank MF085563) was derived from template G1255347364 (GenBank FJ709030, *Peltigera degenii*), the GenBank sequence closest to the new species, with 93% identity for ITS2 region (compared with only 90% identity between the new species and *P. membranacea*). The calculated free energy is -17.80 kcal/mol. Percentages of helix transfer are /100/100/96/85/ (threshold:75%; E-value: $1.2e-52$) for four helices. The computed structure is shown in FIG. 2. One obvious difference between the structural models representing our new species and the closely related *P. degenii* includes one hCBC in the conserved part of helix III and one hCBC and two CBCs in the conserved part of helix IV.

Taxonomy

Peltigera neodegenii L.F. Han, S.Y. Guo & Xiao M. Xu, **sp. nov.**

FIG. 3

MYCOBANK MB821289

Differs from *Peltigera degenii* by its upper surface that is shiny at the centre but tomentose towards the margin, and by the absence of marginal phyllidia.

TYPE: China, Hubei Province, Shennongjia, Jinhouling, 31°28'N 110°18'E, with mosses over rock and soil, 2600 m alt., 24 April 2014, Shou-Yu Guo, Liu-Fu Han 21112. (Holotype, HMAS-L; GenBank MF085563; isotype, HBNU).

ETYMOLOGY: Referring to the morphological similarity with *Peltigera degenii*.

THALLUS foliose, loosely appressed, medium to large, 8–15 cm in diam.; lobes thin, somewhat brittle, 8–20 mm wide, separate or sometimes loosely overlapping, short to elongate, irregularly branching; lobe tips rounded, predominantly downturned; LOBE MARGINS mostly even or weakly crisped, occasionally with lobules; UPPER SURFACE deep bluish green to blackish green (moist) or pale bluish grey to brown (dry), smooth and somewhat shiny centrally, white-gray tomentose near lobe tips, the tomentum becoming absent toward thallus centre, appressed to occasionally erect in part; soredia and isidia lacking, epruinose; LOWER SURFACE distinctly veined; veins usually whitish, not appearing to overlap, narrow, often raised, usually lacking tomentum, but sometimes distinctly erect-tomentose; interstices white, lenticular to polygonal, moderately deeply set; RHIZINES concolourous with veins to pale brown, irregular, discrete, threadlike, occasionally branched, 4.0–8.0 mm long, lacking tomentum or sometimes distinctly erect-tomentose; cortex c. 30–55 µm thick; photobiont layer 25–40 µm thick, containing *Nostoc*; MEDULLA white, c. 250–700 µm thick.

APOTHECIA growing at short lobe tips, erect, saddle-shaped, ≤5.0 mm in diam; apothecial margin smooth to crenulate; DISC red brown to dark brown, smooth; PARAPHYSES simple, septate, ± swollen at the apices and pigmented; ASCI clavate, *Peltigera*-type, colorless to pale, 8-spored; ASCOSPORES acicular, 3(–4)-septate, 60–95 × 2.5–5.5 µm. PYCNIDIA not seen.

SPOT TESTS: all negative.

SECONDARY METABOLITES: none detected.

ECOLOGY & DISTRIBUTION: On mosses and soil in subtropical mountain forest; known only from Shennongjia forest region, Hubei, China.

ADDITIONAL SPECIMENS EXAMINED:

Peltigera neodegenii: CHINA, HUBEI PROVINCE, Shennongjia, Jinhouling, 31°28'N 110°18'E, with mosses over rock and soil, 2600 m alt., 24 April 2014, Shou-Yu Guo, Liu-Fu Han 20973 (HMAS-L, GenBank MF085564); 21105 (HMAS-L, GenBank MF085561); 21111 (HMAS-L, GenBank MF085562).




FIG. 3. *Peltigera neodegenii* (holotype, HMAS-L Guo & Han 21112). a: upper surface and apothecia (dry); b: upper surface, veins, and rhizines (dry); c: section through apothecium (stained by 0.2% toluidine blue for c. 15 min) showing asci and ascospores. Scale bars: a, b = 10 mm; c = 20 µm.

Peltigera praetextata (Flörke ex Sommerf.) Zopf: CHINA, HUBEI PROVINCE, Shennongjia, Laojunshan, 31°49'N 110°32'E, Shou-Yu Guo, Liu-Fu Han 20953 (HMAS-L, GenBank MF085560).

Discussion

The new species shows closer phylogenetic relationships with *Peltigera membranacea* and *P. degenii* than with other species in the ITS sequence analyses (FIG. 1), while the secondary structure models for ITS2 sequences suggest a greater with *P. degenii* (FIG. 2). (In contrast, there was very little similarity with the secondary structure model for the ITS2 sequence of *P. membranacea*, GenBank JX181776 [not shown].)

Morphologically *Peltigera neodegenii* is characterized by the presence of a cyanobacterial photobiont (*Nostoc* sp.) combined with a centrally shiny but marginally tomentose upper surface, predominantly downturned lobe tips, mostly even lobe margins, simple rhizines, and saddle-shaped apothecia. *Peltigera neodegenii* may be confused with *P. membranacea*, which also has a shiny central upper surface shiny and tomentose lobe ends, but which differs by sometimes wider (<4.0 cm) lobes, tomentose veins, and rope-like, tomentose rhizines (Wu & Liu 2012).

Peltigera canina (L.) Willd. and *P. degenii* also resemble *P. neodegenii* in morphology, but *P. canina* differs in its penicillate to confluent rhizines, while *P. degenii* differs in having glossy, glabrous upper surface and distinct marginal phyllidia (Goward & al. 1995; Wu & Liu 2012). *Peltigera evansiana* Gyeln. and *P. praetextata* are readily distinguished from *P. neodegenii*: *P. evansiana* by having granular isidia covering the lobe surfaces, and *P. praetextata* by having phyllidia produced at the margin and along cracks.

Previous studies that have used DNA sequence data for species recognition in *Peltigera* required monophyly both in single-locus ITS phylogenies and diagnosable morphological differences (Goffinet & Miądlikowska 1999; Goward & Goffinet 2000; Goffinet & al. 2003; Miądlikowska & al. 2003; Han & al. 2013, 2015). Here we add secondary structure models for ITS2 sequences, which are also useful characters to distinguish lichen species (Liu & Guo 2009, Cao & al. 2011, Guo 2013, Han & al. 2015). Coleman (2007) considered two organisms as representing different species if the conserved parts of their ITS2 sequences differed by more than one compensatory base change (CBC) or more than four hemi-compensatory base changes (hCBC). The threshold range between intraspecific and interspecific taxa for at least one CBC or four hCBCs in the four helices (I-IV) (Schultz & al. 2005, Guo 2013, Han & al. 2015). For *Peltigera neodegenii* and its template *P. degenii*, there are two hCBCs and two CBCs in the conserved part of helix III and helix IV, further supporting *P. neodegenii* as an independent species.

In conclusion, both distinctive morphological and molecular characteristics of *Peltigera neodegenii* confirm it as new to science.

Acknowledgments

The authors thank Dr. Yoshihito Ohmura (National Museum of Nature and Science, Tsukuba, Ibaraki, Japan) and Dr. Manrong Huang (Beijing Museum of Natural History, Beijing, China) for presubmission review. Prof. Dr. Orvo Vitikainen

(Botanical Museum, Finnish Museum of Natural History, Helsinki) is gratefully acknowledged for providing literature and comments. This study was supported by the Key Technologies R&D Program of China (2013BAD03B03) and the Key Laboratory Open Foundation of Hubei Province (2012snjAB001). Additionally, S.Y. Guo received grant 31370067 from the National Natural Science Foundation of China, and L.F. Han received the grant C2016205201 from the Natural Science Foundation of Hebei Province.

Literature cited

- Byun Y, Han K. 2009. Pseudo Viewer 3: generating planar drawings of large-scale RNA structures with pseudoknots. *Bioinformatics* 25: 1435–1437. <https://doi.org/10.1093/bioinformatics/btp252>
- Cao SN, Liu M, Zhou QM, Guo SY. 2011. Group I introns in lichen forming fungi and their application for phylogenetic analysis. *Mycosystema* 30: 920–931.
- Chen XL. 1986. *Peltigera* in North East China. *Acta Mycologica Sinica* 5: 18–29.
- Coleman AW. 2007. Pan-eukaryote ITS2 homologies revealed by RNA secondary structure. *Nucleic Acids Research* 35: 3322–3329. <https://doi.org/10.1093/nar/gkm233>
- Goffinet B, Miadlikowska J. 1999. *Peltigera phyllidiosa* (*Peltigeraceae*, *Ascomycotina*), a new species from the Southern Appalachians corroborated by ITS sequences. *Lichenologist* 31: 247–256. <https://doi.org/10.1006/lich.1998.0201>
- Goffinet B, Miadlikowska J, Goward T. 2003. Phylogenetic inferences based on nrDNA sequences support five morphospecies within the *Peltigera didactyla* complex (lichenized *Ascomycota*). *Bryologist* 106: 349–364. <https://doi.org/10.1639/01>
- Goward T, Goffinet B. 2000. *Peltigera chionophila*, a new lichen (*Ascomycetes*) from the Western Cordillera of North America. *Bryologist* 103: 493–498. [https://doi.org/10.1639/0007-2745\(2000\)103](https://doi.org/10.1639/0007-2745(2000)103)
- Goward T, Goffinet B, Vitikainen O. 1995. Synopsis of the genus *Peltigera* (lichenized *Ascomycetes*) in British Columbia, with a key to the North American species. *Canadian Journal of Botany* 73: 91–111. <https://doi.org/10.1139/b95-012>
- Guo SY. 2013. Distinguishing species in the lichen genera *Cetrelia* and *Platismatia* (*Ascomycota*) by the new genetic distances inferred from nrDNA ITS sequences. *Chenia* 11: 183–195.
- Han LF, Zhang YY, Guo SY. 2013. *Peltigera wulingensis*, a new lichen (*Ascomycota*) from north China. *Lichenologist* 45: 329–336. <https://dx.doi.org/10.1017/S0024282912000837>
- Han LF, Zheng TX, Guo SY. 2015. A new species in the lichen genus *Peltigera* from northern China based on morphology and DNA sequence data. *Bryologist* 118: 46–53. <https://doi.org/10.1639/0007-2745-118.1.046>
- Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogeny trees. *Bioinformatics* 17: 754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- Kirk PM, Cannon PF, Minter DW, Stalpers JA (eds). 2008. *Dictionary of the Fungi*, 10th edition. CABI Publishing, UK.
- Koetschan C, Förster F, Keller A, Schleicher T, Ruderisch B, Schwarz R, Müller T, Wolf M, Schultz J. 2010. The ITS2 Database III - sequences and structures for phylogeny. *Nucleic Acids Research* 38(Suppl. 1): D275–D279. <https://doi.org/10.1093/nar/gkp966>
- Liu CY, Guo SY. 2009. Comparative analysis of secondary structure of 5.8S-ITS2 rRNA in the genus *Usnea*. *Mycosystema* 28: 705–711.
- Lutzoni F, Wagner P, Reeb V, Zoller S. 2000. Integrating ambiguously aligned regions of DNA sequences in phylogenetic analyses without violating positional homology. *Systematic Biology* 49: 628–651. <https://doi.org/10.1080/106351500750049743>

- Manoharan-Basil SS, Miądlikowska J, Goward T, Andrésson ÓS, Miao VPW. 2016. *Peltigera islandica*, a new cyanolichen species in section *Peltigera* ('*P. canina* group'). *Lichenologist* 48: 451–467. <https://doi.org/10.1017/S0024282916000414>
- Martínez I, Buggazo AR, Vitikainen O, Escudero A. 2003. Distribution patterns in the genus *Peltigera* Willd. *Lichenologist* 35: 301–323. [https://doi.org/10.1016/S0024-2829\(03\)00041-0](https://doi.org/10.1016/S0024-2829(03)00041-0)
- Miądlikowska J, Lutzoni F. 2000. Phylogenetic revision of the genus *Peltigera* (lichen-forming *Ascomycota*) based on morphological, chemical, and large subunit nuclear ribosomal DNA data. *International Journal of Plant Sciences* 161: 925–958. <https://doi.org/10.1086/317568>
- Miądlikowska J, Lutzoni F, Goward T, Zoller S, Posada D. 2003. New approach to an old problem: incorporating signal from gap-rich regions of ITS and rDNA large subunit into phylogenetic analyses to resolve the *Peltigera canina* species complex. *Mycologia* 95: 1181–1203. <https://doi.org/10.2307/3761919>
- Niu D, Zhu Q, Wang Z, Shi J, Bai M, Zheng X. 2016. New record of the lichen genus *Peltigera* from China. *Acta Botanica Boreali-Occidentalia Sinica* 36: 1245–1249. <https://doi.org/10.7606/j.issn.1000-4025.2016.06.1245>
- O'Brien HE, Miądlikowska J, Lutzoni F. 2009. Assessing reproductive isolation in highly diverse communities of the lichen-forming fungal genus *Peltigera*. *Evolution* 63: 2076–2086. <https://doi.org/10.1111/j.1558-5646.2009.00685.x>
- Orange A, James PW, White FJ. 2010. *Microchemical methods for the identification of lichens*, 2nd edition. London: British Lichen Society.
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, & al. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. *Proceedings of the National Academy of Sciences of the United States of America* 109: 6241–6246. <https://doi.org/10.1073/pnas.1117018109>
- Schultz J, Maisel S, Gerlach D, Müller T, Wolf M. 2005. A common core of secondary structure of the internal transcribed spacer 2 (ITS2) throughout the *Eukaryota*. *RNA* 11: 361–364. <https://doi.org/10.1261/rna.7204505>
- Sérusiaux E, Goffinet B, Miądlikowska J, Vitikainen O. 2009. Taxonomy, phylogeny and biogeography of the lichen genus *Peltigera* in Papua New Guinea. *Fungal Diversity* 38: 185–224.
- Stenroos S, Vitikainen O, Koponen T. 1994. *Cladoniaceae, Peltigeraceae* and other lichens from northwestern Sichuan, China. *Journal of the Hattori Botanical Laboratory* 75: 319–344.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* 30: 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Vitikainen O. 1986. *Peltigera dolichospora*, a new Himalayan-Western Chinese lichen. *Lichenologist* 18: 387–390. <https://doi.org/10.1017/S0024282986000580>
- Vitikainen O. 2004. *Peltigera*. 389–398, in: Nash III TH & al. (eds). *Lichen Flora of the Greater Sonoran Desert Region*, Vol. 2. Lichens Unlimited, Tempe, Arizona.
- Wei JC. 1991. *An enumeration of lichens in China*. International Academic Publishers, Beijing.
- White TJ, Bruns TD, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. 315–322, in: MA Innis & al. (eds). *PCR Protocols: a guide to methods and applications*. San Diego, Academic Press. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wu JN, Liu HJ. 2012. *Flora Lichenun Sinicorum*, Vol. 11. Science Press, Beijing.